

10 mL

Must be mixed with
40 mL human milk

Humavant® +4

Human Milk Fortifier (Human, Pasteurised)

Food for special medical purpose

For the dietary management of premature/low-birth-weight infants fed human milk. Product must be used under medical supervision. Not for parenteral use.

Product Description

Humavant® +4 human milk fortifier (HMF) is the first HMF made exclusively from 100% donor breast milk. The product is a human milk-based, concentrated, pasteurised, liquid HMF that helps provide essential calories, protein, and minerals to meet the nutritional needs of premature infants.

- Nutritionally incomplete. Infant will require additional vitamins and iron added separately from the product.
- Available frozen in 125 mL bottles containing 10 mL of product.

Ingredients

Human milk, calcium glycerophosphate, sodium citrate, potassium citrate, calcium chloride, calcium gluconate, magnesium phosphate, sodium chloride, zinc sulfate, cupric sulfate.

Storage

Store at -20°C or colder until ready to thaw for use.

Directions for Thawing

Under no circumstances should the product be defrosted or warmed in a microwave.

Recommended method of thawing is refrigeration (2°C to 8°C).

- Place unopened (frozen) bottle in refrigerator for 2-5 hours. Thaw time will vary by fortifier concentration. As the caloric value (volume) increases, the thaw time may take longer.
- Swirl gently to detect ice in the bottle. If ice is still present, return to the refrigerator for additional thaw time. Repeat until no ice is detected.
- Once the thawing process begins, administer within 48 hours.
- Do not refreeze. Keep refrigerated until used.

Preparation Instructions

Always maintain aseptic technique when preparing and handling human milk. DO NOT ADD WATER.

Each bottle of Humavant +4 fortifier contains 10 mL of fortifier and must be mixed with 40 mL of human milk (ratio 1:4).

- After the bottle has been properly thawed (see above), remove the cap from bottle.
- Add 40 mL of human milk (expressed breast milk or donor milk) into the Humavant +4 bottle to achieve a total volume of 50 mL.
- Gently swirl bottle to mix; DO NOT SHAKE. The product is now ready for use.
- Measure out the fortified milk using sterile syringes according to the feeding order.
- Label each syringe with patient identifier and refrigerate (2°C to 8°C) until administered.

References:

1. American Academy of Pediatrics Committee on Nutrition. Appendix A. In: Kleinman RE, Greer FR, eds. *Pediatric Nutrition*. 8th ed. Itasca, IL: American Academy of Pediatrics; 2019:1505-1508.
2. Mimouni FB, Mandel D, Lubetzky R, Senterre T. Calcium, phosphorus, magnesium and vitamin D requirements of the preterm infant. *World Rev Nutr Diet*. 2014;110:140-151. doi:10.1159/000358463

Use of Product

Initiation of enteral feedings and advancement rates should be individualized based on infant's weight, age, and clinical status. Typically, mother's own milk or donor human milk provides 65-70 kcal (272-293 kJ)/100 mL.¹ Calcium to phosphorus ratio supports intrauterine accretion rates.² When used under medical supervision, both protein and fat levels are appropriate to achieve adequate growth.

When mixed with 0.67 kcal (3 kJ) / mL preterm human milk, Humavant +4 provides 42 kcal (176 kJ) and 1.3 grams of protein per 50 mL of feeding solution. Nutrition monitoring is always required.

An Exclusive Human Milk Diet (EHMD)

An EHMD is achieved when 100% of the protein, fat, and carbohydrates are derived solely from human milk. Humavant + fortifier is the first HMF derived from human milk as opposed to cow milk.

Prolacta Bioscience offers a full line of human milk-based products for providing an EHMD. If mother's own milk cannot be assured to provide a minimum of 0.67 kcal (3 kJ) / mL, Humavant® CR human milk caloric fortifier can be used. Humavant CR fortifier is a pasteurised formulation of human milk cream (derived from donor human milk) that can be added to mother's milk to increase the caloric content to 0.67 kcal (3 kJ) / mL.

An EHMD may require additional nutrients. No commercially-available HMF can be guaranteed to provide the full and necessary nutritional needs of every preterm infant.

Safety Information

Abruptly transitioning the infant's diet from this product to cow milk-based nutrition could result in feeding intolerance or gastrointestinal complications. To obtain a copy of Prolacta Bioscience's *Feeding Transition From an Exclusive Human Milk Diet*, please contact your Prolacta Bioscience Representative.

Distributed By

Prolacta Bioscience Australia PTY LTD
Ground Floor
121 King William Street, Adelaide
South Australia, 5000
1 800955840
Prolacta.com/as

MS-0136



20 mL

Must be mixed with
80 mL human milk

Humavant® +4

Human Milk Fortifier (Human, Pasteurised)

Food for special medical purpose

For the dietary management of premature/low-birth-weight infants fed human milk. Product must be used under medical supervision. Not for parenteral use.

Product Description

Humavant® +4 human milk fortifier (HMF) is the first HMF made exclusively from 100% donor breast milk. The product is a human milk-based, concentrated, pasteurised, liquid HMF that helps provide essential calories, protein, and minerals to meet the nutritional needs of premature infants.

- Nutritionally incomplete. Infant will require additional vitamins and iron added separately from the product.
- Available frozen in 125 mL bottles containing 20 mL of product.

Ingredients

Human milk, calcium glycerophosphate, sodium citrate, potassium citrate, calcium chloride, calcium gluconate, magnesium phosphate, sodium chloride, zinc sulfate, cupric sulfate.

Storage

Store at -20°C or colder until ready to thaw for use.

Directions for Thawing

Under no circumstances should the product be defrosted or warmed in a microwave.

Recommended method of thawing is refrigeration (2°C to 8°C).

- Place unopened (frozen) bottle in refrigerator for 2-5 hours. Thaw time will vary by fortifier concentration. As the caloric value (volume) increases, the thaw time may take longer.
- Swirl gently to detect ice in the bottle. If ice is still present, return to the refrigerator for additional thaw time. Repeat until no ice is detected.
- Once the thawing process begins, administer within 48 hours.
- Do not refreeze. Keep refrigerated until used.

Preparation Instructions

Always maintain aseptic technique when preparing and handling human milk. DO NOT ADD WATER.

Each bottle of Humavant +4 fortifier contains 20 mL of fortifier and must be mixed with 80 mL of human milk (ratio 1:4).

- After the bottle has been properly thawed (see above), remove the cap from bottle.
- Add 80 mL of human milk (expressed breast milk or donor milk) into the Humavant +4 bottle to achieve a total volume of 100 mL.
- Gently swirl bottle to mix; DO NOT SHAKE. The product is now ready for use.
- Measure out the fortified milk using sterile syringes according to the feeding order.
- Label each syringe with patient identifier and refrigerate (2°C to 8°C) until administered.

References:

1. American Academy of Pediatrics Committee on Nutrition. Appendix A. In: Kleinman RE, Greer FR, eds. *Pediatric Nutrition*. 8th ed. Itasca, IL: American Academy of Pediatrics; 2019:1505-1508.
2. Mimouni FB, Mandel D, Lubetzky R, Senterre T. Calcium, phosphorus, magnesium and vitamin D requirements of the preterm infant. *World Rev Nutr Diet*. 2014;110:140-151. doi:10.1159/000358463

Use of Product

Initiation of enteral feedings and advancement rates should be individualized based on infant's weight, age, and clinical status. Typically, mother's own milk or donor human milk provides 65-70 kcal (272-293 kJ)/100 mL.¹ Calcium to phosphorus ratio supports intrauterine accretion rates.² When used under medical supervision, both protein and fat levels are appropriate to achieve adequate growth.

When mixed with 0.67 kcal (3 kJ) / mL preterm human milk, Humavant +4 provides 83 kcal (347 kJ) and 2.5 grams of protein per 100 mL of feeding solution. Nutrition monitoring is always required.

An Exclusive Human Milk Diet (EHMD)

An EHMD is achieved when 100% of the protein, fat, and carbohydrates are derived solely from human milk. Humavant + fortifier is the first HMF derived from human milk as opposed to cow milk.

Prolacta Bioscience offers a full line of human milk-based products for providing an EHMD. If mother's own milk cannot be assured to provide a minimum of 0.67 kcal (3 kJ) / mL, Humavant® CR human milk caloric fortifier can be used. Humavant CR fortifier is a pasteurised formulation of human milk cream (derived from donor human milk) that can be added to mother's milk to increase the caloric content to 0.67 kcal (3 kJ) / mL.

An EHMD may require additional nutrients. No commercially-available HMF can be guaranteed to provide the full and necessary nutritional needs of every preterm infant.

Safety Information

Abruptly transitioning the infant's diet from this product to cow milk-based nutrition could result in feeding intolerance or gastrointestinal complications. To obtain a copy of Prolacta Bioscience's *Feeding Transition From an Exclusive Human Milk Diet*, please contact your Prolacta Bioscience Representative.

Distributed By

Prolacta Bioscience Australia PTY LTD
Ground Floor
121 King William Street, Adelaide
South Australia, 5000
1 800955840
Prolacta.com/as

MS-0137



15 mL

Must be mixed with
35 mL human milk

Humavant[®] +6

Human Milk Fortifier (Human, Pasteurised)

Food for special medical purpose

For the dietary management of premature/low-birth-weight infants fed human milk. Product must be used under medical supervision. Not for parenteral use.

Product Description

Humavant[®] +6 human milk fortifier (HMF) is the first HMF made exclusively from 100% donor breast milk. The product is a human milk-based, concentrated, pasteurised, liquid HMF that helps provide essential calories, protein, and minerals to meet the nutritional needs of premature infants.

- Nutritionally incomplete. Infant will require additional vitamins and iron added separately from the product.
- Available frozen in 125 mL bottles containing 15 mL of product.

Ingredients

Human milk, calcium glycerophosphate, sodium citrate, potassium citrate, calcium chloride, calcium gluconate, magnesium phosphate, sodium chloride, zinc sulfate, cupric sulfate.

Storage

Store at -20°C or colder until ready to thaw for use.

Directions for Thawing

Under no circumstances should the product be defrosted or warmed in a microwave.

Recommended method of thawing is refrigeration (2°C to 8°C).

- Place unopened (frozen) bottle in refrigerator for 2-5 hours. Thaw time will vary by fortifier concentration. As the caloric value (volume) increases, the thaw time may take longer.
- Swirl gently to detect ice in the bottle. If ice is still present, return to the refrigerator for additional thaw time. Repeat until no ice is detected.
- Once the thawing process begins, administer within 48 hours.
- Do not refreeze. Keep refrigerated until used.

Preparation Instructions

Always maintain aseptic technique when preparing and handling human milk. DO NOT ADD WATER.

Each bottle of Humavant +6 fortifier contains 15 mL of fortifier and must be mixed with 35 mL of human milk (ratio 3:7).

- After the bottle has been properly thawed (see above), remove the cap from bottle.
- Add 35 mL of human milk (expressed breast milk or donor milk) into the Humavant +6 bottle to achieve a total volume of 50 mL.
- Gently swirl bottle to mix; DO NOT SHAKE. The product is now ready for use.
- Measure out the fortified milk using sterile syringes according to the feeding order.
- Label each syringe with patient identifier and refrigerate (2°C to 8°C) until administered.

References:

1. American Academy of Pediatrics Committee on Nutrition. Appendix A. In: Kleinman RE, Greer FR, eds. *Pediatric Nutrition*. 8th ed. Itasca, IL: American Academy of Pediatrics; 2019:1505-1508.
2. Mimouni FB, Mandel D, Lubetzky R, Senterre T. Calcium, phosphorus, magnesium and vitamin D requirements of the preterm infant. *World Rev Nutr Diet*. 2014;110:140-151. doi:10.1159/000358463

Use of Product

Initiation of enteral feedings and advancement rates should be individualized based on infant's weight, age, and clinical status. Typically, mother's own milk or donor human milk provides 65-70 kcal (272-293 kJ)/100 mL.¹ Calcium to phosphorus ratio supports intrauterine accretion rates.² When used under medical supervision, both protein and fat levels are appropriate to achieve adequate growth.

When mixed with 0.67 kcal (3 kJ) / mL preterm human milk, Humavant +6 provides 45 kcal (188 kJ) and 1.5 grams of protein per 50 mL of feeding solution. Nutrition monitoring is always required.

An Exclusive Human Milk Diet (EHMD)

An EHMD is achieved when 100% of the protein, fat, and carbohydrates are derived solely from human milk. Humavant + fortifier is the first HMF derived from human milk as opposed to cow milk.

Prolacta Bioscience offers a full line of human milk-based products for providing an EHMD. If mother's own milk cannot be assured to provide a minimum of 0.67 kcal (3 kJ) / mL, Humavant[®] CR human milk caloric fortifier can be used. Humavant CR fortifier is a pasteurised formulation of human milk cream (derived from donor human milk) that can be added to mother's milk to increase the caloric content to 0.67 kcal (3 kJ) / mL.

An EHMD may require additional nutrients. No commercially-available HMF can be guaranteed to provide the full and necessary nutritional needs of every preterm infant.

Safety Information

Abruptly transitioning the infant's diet from this product to cow milk-based nutrition could result in feeding intolerance or gastrointestinal complications. To obtain a copy of Prolacta Bioscience's *Feeding Transition From an Exclusive Human Milk Diet*, please contact your Prolacta Bioscience Representative.

Distributed By

Prolacta Bioscience Australia PTY LTD
Ground Floor
121 King William Street, Adelaide
South Australia, 5000
1 800955840
Prolacta.com/as

MS-0138



30 mL

Must be mixed with
70 mL human milk

Humavant® +6

Human Milk Fortifier (Human, Pasteurised)

Food for special medical purpose

For the dietary management of premature/low-birth-weight infants fed human milk. Product must be used under medical supervision. Not for parenteral use.

Product Description

Humavant® +6 human milk fortifier (HMF) is the first HMF made exclusively from 100% donor breast milk. The product is a human milk-based, concentrated, pasteurised, liquid HMF that helps provide essential calories, protein, and minerals to meet the nutritional needs of premature infants.

- Nutritionally incomplete. Infant will require additional vitamins and iron added separately from the product.
- Available frozen in 125 mL bottles containing 30 mL of product.

Ingredients

Human milk, calcium glycerophosphate, sodium citrate, potassium citrate, calcium chloride, calcium gluconate, magnesium phosphate, sodium chloride, zinc sulfate, cupric sulfate.

Storage

Store at -20°C or colder until ready to thaw for use.

Directions for Thawing

Under no circumstances should the product be defrosted or warmed in a microwave.

Recommended method of thawing is refrigeration (2°C to 8°C).

- Place unopened (frozen) bottle in refrigerator for 2-5 hours. Thaw time will vary by fortifier concentration. As the caloric value (volume) increases, the thaw time may take longer.
- Swirl gently to detect ice in the bottle. If ice is still present, return to the refrigerator for additional thaw time. Repeat until no ice is detected.
- Once the thawing process begins, administer within 48 hours.
- Do not refreeze. Keep refrigerated until used.

Preparation Instructions

Always maintain aseptic technique when preparing and handling human milk. DO NOT ADD WATER.

Each bottle of Humavant +6 fortifier contains 30 mL of fortifier and must be mixed with 70 mL of human milk (ratio 3:7).

- After the bottle has been properly thawed (see above), remove the cap from bottle.
- Add 70 mL of human milk (expressed breast milk or donor milk) into the Humavant +6 bottle to achieve a total volume of 100 mL.
- Gently swirl bottle to mix; DO NOT SHAKE. The product is now ready for use.
- Measure out the fortified milk using sterile syringes according to the feeding order.
- Label each syringe with patient identifier and refrigerate (2°C to 8°C) until administered.

References:

1. American Academy of Pediatrics Committee on Nutrition. Appendix A. In: Kleinman RE, Greer FR, eds. *Pediatric Nutrition*. 8th ed. Itasca, IL: American Academy of Pediatrics; 2019:1505-1508.
2. Mimouni FB, Mandel D, Lubetzky R, Senterre T. Calcium, phosphorus, magnesium and vitamin D requirements of the preterm infant. *World Rev Nutr Diet*. 2014;110:140-151. doi:10.1159/000358463

Use of Product

Initiation of enteral feedings and advancement rates should be individualized based on infant's weight, age, and clinical status. Typically, mother's own milk or donor human milk provides 65-70 kcal (272-293 kJ)/100 mL.¹ Calcium to phosphorus ratio supports intrauterine accretion rates.² When used under medical supervision, both protein and fat levels are appropriate to achieve adequate growth.

When mixed with 0.67 kcal (3 kJ) / mL preterm human milk, Humavant +6 provides 91 kcal (381 kJ) and 2.9 grams of protein per 100 mL of feeding solution. Nutrition monitoring is always required.

An Exclusive Human Milk Diet (EHMD)

An EHMD is achieved when 100% of the protein, fat, and carbohydrates are derived solely from human milk. Humavant + fortifier is the first HMF derived from human milk as opposed to cow milk.

Prolacta Bioscience offers a full line of human milk-based products for providing an EHMD. If mother's own milk cannot be assured to provide a minimum of 0.67 kcal (3 kJ) / mL, Humavant® CR human milk caloric fortifier can be used. Humavant CR fortifier is a pasteurised formulation of human milk cream (derived from donor human milk) that can be added to mother's milk to increase the caloric content to 0.67 kcal (3 kJ) / mL.

An EHMD may require additional nutrients. No commercially-available HMF can be guaranteed to provide the full and necessary nutritional needs of every preterm infant.

Safety Information

Abruptly transitioning the infant's diet from this product to cow milk-based nutrition could result in feeding intolerance or gastrointestinal complications. To obtain a copy of Prolacta Bioscience's *Feeding Transition From an Exclusive Human Milk Diet*, please contact your Prolacta Bioscience Representative.

Distributed By

Prolacta Bioscience Australia PTY LTD
Ground Floor
121 King William Street, Adelaide
South Australia, 5000
1 800955840
Prolacta.com/as

MS-0139



40 mL

Must be mixed with
60 mL human milk

Humavant[®] +8

Human Milk Fortifier (Human, Pasteurised)

Food for special medical purpose

For the dietary management of premature/low-birth-weight infants fed human milk. Product must be used under medical supervision. Not for parenteral use.

Product Description

Humavant[®] +8 human milk fortifier (HMF) is the first HMF made exclusively from 100% donor breast milk. The product is a human milk-based, concentrated, pasteurised, liquid HMF that helps provide essential calories, protein, and minerals to meet the nutritional needs of premature infants.

- Nutritionally incomplete. Infant will require additional vitamins and iron added separately from the product.
- Available frozen in 125 mL bottles containing 40 mL of product.

Ingredients

Human milk, calcium glycerophosphate, potassium citrate, sodium citrate, calcium chloride, calcium gluconate, sodium chloride, magnesium phosphate, zinc sulfate, cupric sulfate.

Storage

Store at -20°C or colder until ready to thaw for use.

Directions for Thawing

Under no circumstances should the product be defrosted or warmed in a microwave.

Recommended method of thawing is refrigeration (2°C to 8°C).

- Place unopened (frozen) bottle in refrigerator for 2-5 hours. Thaw time will vary by fortifier concentration. As the caloric value (volume) increases, the thaw time may take longer.
- Swirl gently to detect ice in the bottle. If ice is still present, return to the refrigerator for additional thaw time. Repeat until no ice is detected.
- Once the thawing process begins, administer within 48 hours.
- Do not refreeze. Keep refrigerated until used.

Preparation Instructions

Always maintain aseptic technique when preparing and handling human milk. DO NOT ADD WATER.

Each bottle of Humavant +8 fortifier contains 40 mL of fortifier and must be mixed with 60 mL of human milk (ratio 2:3).

- After the bottle has been properly thawed (see above), remove the cap from bottle.
- Add 60 mL of human milk (expressed breast milk or donor milk) into the Humavant +8 bottle to achieve a total volume of 100 mL.
- Gently swirl bottle to mix; DO NOT SHAKE. The product is now ready for use.
- Measure out the fortified milk using sterile syringes according to the feeding order.
- Label each syringe with patient identifier and refrigerate (2°C to 8°C) until administered.

References:

1. American Academy of Pediatrics Committee on Nutrition. Appendix A. In: Kleinman RE, Greer FR, eds. *Pediatric Nutrition*. 8th ed. Itasca, IL: American Academy of Pediatrics; 2019:1505-1508.
2. Mimouni FB, Mandel D, Lubetzky R, Senterre T. Calcium, phosphorus, magnesium and vitamin D requirements of the preterm infant. *World Rev Nutr Diet*. 2014;110:140-151. doi:10.1159/000358463

Use of Product

Initiation of enteral feedings and advancement rates should be individualized based on infant's weight, age, and clinical status. Typically, mother's own milk or donor human milk provides 65-70 kcal (272-293 kJ)/100 mL.¹ Calcium to phosphorus ratio supports intrauterine accretion rates.² When used under medical supervision, both protein and fat levels are appropriate to achieve adequate growth.

When mixed with 0.67 kcal (3 kJ) / mL preterm human milk, Humavant +8 provides 98 kcal (410 kJ) and 3.4 grams of protein per 100 mL of feeding solution. Extreme caution should be taken when using Humavant +8 fortified milk at volumes above 150 mL/kg/day due to high protein intake. Nutrition monitoring is always required.

An Exclusive Human Milk Diet (EHMD)

An EHMD is achieved when 100% of the protein, fat, and carbohydrates are derived solely from human milk. Humavant + fortifier is the first HMF derived from human milk as opposed to cow milk.

Prolacta Bioscience offers a full line of human milk-based products for providing an EHMD. If mother's own milk cannot be assured to provide a minimum of 0.67 kcal (3 kJ) / mL, Humavant[®] CR human milk caloric fortifier can be used. Humavant CR fortifier is a pasteurised formulation of human milk cream (derived from donor human milk) that can be added to mother's milk to increase the caloric content to 0.67 kcal (3 kJ) / mL.

An EHMD may require additional nutrients. No commercially-available HMF can be guaranteed to provide the full and necessary nutritional needs of every preterm infant.

Safety Information

Abruptly transitioning the infant's diet from this product to cow milk-based nutrition could result in feeding intolerance or gastrointestinal complications. To obtain a copy of Prolacta Bioscience's *Feeding Transition From an Exclusive Human Milk Diet*, please contact your Prolacta Bioscience Representative.

Distributed By

Prolacta Bioscience Australia PTY LTD
Ground Floor
121 King William Street, Adelaide
South Australia, 5000
1 800955840
Prolacta.com/as

MS-0140



50 mL

Must be mixed with
50 mL human milk

Humavant® +10

Human Milk Fortifier (Human, Pasteurised)

Food for special medical purpose

For the dietary management of premature/low-birth-weight infants fed human milk. Product must be used under medical supervision. Not for parenteral use.

Product Description

Humavant® +10 human milk fortifier (HMF) is the first HMF made exclusively from 100% donor breast milk. The product is a human milk-based, concentrated, pasteurised, liquid HMF that helps provide essential calories, protein, and minerals to meet the nutritional needs of premature infants.

- Nutritionally incomplete. Infant will require additional vitamins and iron added separately from the product.
- Available frozen in 125 mL bottles containing 50 mL of product.

Ingredients

Human milk, calcium glycerophosphate, potassium citrate, sodium citrate, calcium chloride, calcium gluconate, sodium chloride, magnesium phosphate, zinc sulfate, cupric sulfate.

Storage

Store at -20°C or colder until ready to thaw for use.

Directions for Thawing

Under no circumstances should the product be defrosted or warmed in a microwave.

Recommended method of thawing is refrigeration (2°C to 8°C).

- Place unopened (frozen) bottle in refrigerator for 2-5 hours. Thaw time will vary by fortifier concentration. As the caloric value (volume) increases, the thaw time may take longer.
- Swirl gently to detect ice in the bottle. If ice is still present, return to the refrigerator for additional thaw time. Repeat until no ice is detected.
- Once the thawing process begins, administer within 48 hours.
- Do not refreeze. Keep refrigerated until used.

Preparation Instructions

Always maintain aseptic technique when preparing and handling human milk. DO NOT ADD WATER.

Each bottle of Humavant +10 fortifier contains 50 mL of fortifier and must be mixed with 50 mL of human milk (ratio 1:1).

- After the bottle has been properly thawed (see above), remove the cap from bottle.
- Add 50 mL of human milk (expressed breast milk or donor milk) into the Humavant +10 bottle to achieve a total volume of 100 mL.
- Gently swirl bottle to mix; DO NOT SHAKE. The product is now ready for use.
- Measure out the fortified milk using sterile syringes according to the feeding order.
- Label each syringe with patient identifier and refrigerate (2°C to 8°C) until administered.

References:

1. American Academy of Pediatrics Committee on Nutrition. Appendix A. In: Kleinman RE, Greer FR, eds. *Pediatric Nutrition*. 8th ed. Itasca, IL: American Academy of Pediatrics; 2019:1505-1508.
2. Mimouni FB, Mandel D, Lubetzky R, Senterre T. Calcium, phosphorus, magnesium and vitamin D requirements of the preterm infant. *World Rev Nutr Diet*. 2014;110:140-151. doi:10.1159/000358463

Use of Product

Initiation of enteral feedings and advancement rates should be individualized based on infant's weight, age, and clinical status. Typically, mother's own milk or donor human milk provides 65-70 kcal (272-293 kJ)/100 mL.¹ Calcium to phosphorus ratio supports intrauterine accretion rates.² When used under medical supervision, both protein and fat levels are appropriate to achieve adequate growth.

When mixed with 0.67 kcal (3 kJ) / mL preterm human milk, Humavant+10 provides 105 kcal (439 kJ) and 3.8 grams of protein per 100 mL of feeding solution. Extreme caution should be taken when using Humavant+10 fortified milk at volumes above 150 mL/kg/day due to high protein intake. Nutrition monitoring is always required.

An Exclusive Human Milk Diet (EHMD)

An EHMD is achieved when 100% of the protein, fat, and carbohydrates are derived solely from human milk. Humavant + fortifier is the first HMF derived from human milk as opposed to cow milk.

Prolacta Bioscience offers a full line of human milk-based products for providing an EHMD. If mother's own milk cannot be assured to provide a minimum of 0.67 kcal (3 kJ) / mL, Humavant® CR human milk caloric fortifier can be used. Humavant CR fortifier is a pasteurised formulation of human milk cream (derived from donor human milk) that can be added to mother's milk to increase the caloric content to 0.67 kcal (3 kJ) / mL.

An EHMD may require additional nutrients. No commercially-available HMF can be guaranteed to provide the full and necessary nutritional needs of every preterm infant.

Safety Information

Abruptly transitioning the infant's diet from this product to cow milk-based nutrition could result in feeding intolerance or gastrointestinal complications. To obtain a copy of Prolacta Bioscience's *Feeding Transition From an Exclusive Human Milk Diet*, please contact your Prolacta Bioscience Representative.

Distributed By

Prolacta Bioscience Australia PTY LTD
Ground Floor
121 King William Street, Adelaide
South Australia, 5000
1 800955840
Prolacta.com/as

MS-0141

