Prolact+4 H²MF[®] Human Milk-Based Human Milk Fortifier (20 mL) Product Preparation Log Sheet

This log sheet should be completed by the staff who will prepare Prolact+4 H²MF fortifier (20 mL) for administration and feeding.

| Directions for Preparation and Use Store at -20°C or colder until ready to use. Do not microwave. Once thawing process begins, administer | 20 mL Prolact+4 H ² MF | Caution Nutritionally incomplete. Infant will require additional vitamins and iron added separately from the product. | | 20 mL Aut be mixed with 10 mL herrary milk |
|--|---|--|-------|---|
| within 48 hours. Mix full bottle with 80 mL of human milk. Gently swid bottle to mix: DO NOT SHAKE. Once | Must be mixed with 80 mL human milk | Manufactured By | | |
| milk: Gently swirt bottle to mix; DO NOI SHARE: Once mixed, keep refrigerated until used. The health of the infant depends on carefully following the directions for areparation and use. | Human Milk Fortifier (Human, Pasteurized) When mixed, adds 4 Calories per fl oz. | Profacto Bioscience, Inc. City of Industry, CA 91746 www.prolacta.com 1/8889 IPROJACT | | Human Milk Fortifi When mixed, add |
| Nutrients per 20 mL (1 bottie) Protein | To be used under the supervision of a physician. Week and the supervision of the honors mit. When must, active California are if a con- | Ingredients Human milk. contains less than 2% of the following: calcium glycorophosphate, calcium glucontet, sodium citrate, magnesium citrate, calcium catronite, zins suffate, composition, calcium catronite, zins suffate, calcium catronite, zins suffate, composition, calcium catronite, zins suffate, calcium catronite, calcium catronite, calcium calcium catronite, calcium catronite, calcium calcium catronite, calcium catronite, calcium calcium catronite, calcium catronite, calcium calcium calcium catronite, calcium calcium calcium catronite, calcium ca | Lot # | To be used under the Intended for premoture When mixed, o |

| Patient Name and MR# or Patient Label | Lot/Serial# | Date/Time Frozen Bottle Removed from Freezer | Discard Bottle by Date/Time* | mL Dispensed∕ mL Remaining† | mL Discarded† | Signature of Preparer/ Co-Signer |
|--|-------------|---|---------------------------------|--------------------------------|------------------|--|
| Example: J. Doe ABC123XYZ | CF041802USA | Date: XX/XX/XX | Date: XX/XX/XX | 80 mL | 0 mL | J. Smith |
| | 5951 | Time: 8:00 am | Time: 8:00 am | 20 mL | | S. Jones |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |

*Product must be administered within 48 hours of the start of thawing or according to hospital policy regarding human milk – whichever time frame is shorter. tWhen mixed with 80 mL of human milk for a total volume of 100 mL.

To provide your preterm patient with a 100% human milk-based diet, call 1-888-PROLACT (1-888-776-5228) or visit www.prolacta.com.



olact+4 H²MF

L (0.7 fl oz)