

Prolact+10 H²MF[®] Human Milk-Based Human Milk Fortifier (50 mL) Product Preparation Log Sheet

This log sheet should be completed by the staff who will prepare Prolact+10 H²MF fortifier (50 mL) for administration and feeding.

Directions for Preparation and Use
Store at 20°C or colder until ready to use. Do not microwave. Once thawing process begins, administer within 48 hours. Mix full bottle with 50 mL of human milk. Gently swirl bottle to mix; DO NOT SHAKE. Once mixed, keep refrigerated until used. The health of the infant depends on carefully following the directions for preparation and use.

50 mL Prolact+10 H²MF[®]
Must be mixed with 50 mL human milk

Human Milk Fortifier (Human, Pasteurized)
When mixed, adds 10 Calories per fl oz.

Nutrients per 50 mL (1 bottle)
Protein..... g Carbohydrate..... g
Fat..... g Calories..... g

Minerals
Calcium..... mg Manganese..... mcg
Phosphorus..... mg Copper..... mcg
Magnesium..... mg Sodium..... mg
Iron..... mg Potassium..... mg
Zinc..... mg Chloride..... mg

Ingredients
Human milk, contains less than 2% of the following: calcium glycerophosphate, calcium gluconate, sodium citrate, magnesium phosphate, calcium chloride, potassium citrate, calcium carbonate, zinc sulfate, cupric sulfate.

Caution
Nutritionally incomplete. Infant will require additional vitamins and iron added separately from the product.

Manufactured by
Prolacta Bioscience, Inc.
City of Industry, CA 91746
www.prolacta.com
1(888) PROLACT

To be used under the supervision of a physician. Intended for premature/LBW infants fed human milk. When mixed, adds 10 Calories per fl oz.

Product Code: 93901-50
GTIN: 03812697021050
LOT: XXXXXXXXXXXX
USE BY: Month Day, Year
SERIAL: XXXXX

+10
50 mL (1.7 fl oz)

Lot #



Patient Name and MR# or Patient Label	Lot/Serial#	Date/Time Frozen Bottle Removed from Freezer	Discard Bottle by Date/Time*	mL Dispensed/ mL Remaining [†]	mL Discarded [†]	Signature of Preparer/ Co-Signer
Example: J. Doe ABC123XYZ	CF101801USA	Date: XX/XX/XX	Date: XX/XX/XX	80 mL	0 mL	J. Smith
	5951	Time: 8:00 am	Time: 8:00 am	20 mL		S. Jones

*Product must be administered within 48 hours of the start of thawing or according to hospital policy regarding human milk – whichever time frame is shorter.

[†]When mixed with 50 mL of human milk for a total volume of 100 mL.

To provide your preterm patient with a 100% human milk-based diet, call 1-888-PROLACT (1-888-776-5228) or visit www.prolacta.com.